REFERENCES

1. E. P. Abraham and G. G. F. Newton, Biochem. J., 79 (1961) 377.
2. D. C. Hodgkin and E. N. Maslem, Biochem. J., 79 (1961) 393.
3. G. G. F. Newton and E. P. Abraham, Biochem. J., 62 (1956) 651.
4. Newton GGF and Abraham EP (1915), Nature, 175, 548.
5. L. M. Gerald, Merl A.C. Penicillins, Cephalosporin and other ß-Lactam Antibiotics. Goodman and Gilmans the Pharmacological Basis of Therapeutics New York, Pargamon Press, (1990) p.1065
6. Rolinson GN (1986), J. Antimicrob. Chemother., 17 (5).
7. Roberts J (1992), The cephalosporins in kirk-othemer encyclopedia of chemical Technology, John wiley and sons. lnc.,vo! 3, 28-82.
8. Elks J (1987), Drugs, 34, Suppl. 2, 240-247
9. Newall CH and Hallam PD in sammes PG and Taylor JB, Eds (1992)., comprehensive medicinal chemistry, pergamon, 2, 609-653.
10. Klein NC and Gunha BA(1995), Advances in therapy, 21, 83.
11. Durkheimer W, Adam F., Fischer G, and Kirrstetter, in B. Testa (1998), Ed., Advances in drug research, academic Press, 17,61-234.
12. Okamoto Y, Kiriyama K, Namiki Y, Matsushita J, Fujioka M and Yasuda T. (1996), Degradation kinetics and isomerization of cefdinir, a new oral cephalosporin, in aqueous solution, J Pharm Sci. Sep;85(9):976-83.
13. Okamoto Y, Itoh K, Namiki Y, Matsushita J, Fujioka M, Yasuda T. (1996), Method development for the determination of cefdinir and its related substances by high-performance liquid chromatography, J Pharm Biomed Anal. 1996 Apr;14(6):739-48.
14. Kees F, Wellenhofer M, Grobecker H, (1996), Bioavailability of cefpodoxime proxetil with co-administered acetylcysteine. Arzneimittelforschang. 46(4), 435-438.
15. Camus F, Deslandes A, Harcouet L, Farinotii R, (1994), High Performance liquid chromatographic method for the determination of cefpodoxime levels in plasma and sinus, mucosa. J. Chromatogr B Biorned. Appl, 656(2), 383-388.
16. N. H. Vadia, N. B. Dobaria and V. B. Patel, (2009), Conventional and Advanced Spectrophotometric Methods for estimation of Cefetamet Pivoxil Hydrochloride in Bulk

and in Pharmaceutical Formulation, International Journal of ChemTech Research,Vol.1, No.2, pp 286-290, April-June 2009

17. Madhusudana Reddy T, Shreedhar M and Jayarama Reddy S, (2003), Voltametric behaviour of cefixime and cefpodoxime proxetil and determination in pharmaceutical formulations and urine. J. Pharm. Biomed. Anal., 31(4), 811-818.
18. Silber Michael B, Andrew J. Falkowski, Zee M.Look, Hideyo Noguchi.Determination of cefixime in biological samples by Reverse-Phase High Performance liquid Chromatography .Journal of chromatography.1987; 422: 145-152.
19. Castillo M, V.Girona, C.Pacareu, A.Riera, R.Pouplana, J.Bolos. Spectrophotometric determination of the stability of an ampicillin-dicloxacillin suspension. Journal of Pharmaceutical & Biomedical Analysis. 1988; 6 (1): 23-28.
20. Eric-Jovanovic S, D.Agbaba, D.Zivanov-Stakic, S.Vladimirov. HPTLC determination of ceftriaxone, cefixime and cefotaxime in dosage forms.Journal of Pharmaceutical and Biomedical Analysis. 1998;18: 893-898
21. Gehad G Mohamed, F.A.Nour EL-Dien, Eman U Farag. Spectrophotometric study of the mechanism between DDQ as π-acceptor and potassium iodate and flucloxacillin and dicloxacillin drugs and their determination in pure and in dosage forms. ScienceDirect Spectrochimica Acta Part A. 2006; 65: 11-19.
22. Nanda RK, Gaikwad J, Prakash A. Simultenous spectrophotometric estimation of cefixime and ornidazole in tablet dosage form. International Journal of PharmaTech Research. 2009; 1(3): 488-491.
23. Shahnaz Gauhar, Hafiz Muhammad Arshad, Raheela Bano, Iyad Naeem Muhammad. Development of HPLC-UV Method for Analysis of cefixime in Raw Materials and in Capsules. Jordan Journal of Pharmaceutical Sciences. 2009; 2(1): 53-65.
24. Madhura V. Dhoka. Vandana T. Gawande, Pranav P. Joshi. Simultenous estimation of Cefixime Trihydrate and Erdostein in Pharmaceutical dosage

form by using RP-HPLC. International Journal of ChemTech Research. 2010; 2(1): 79-87.

25. M.K Kumudhavalli, Sandeep Sahu, K. Abhiteja, Dr.B. Jayakar. Development and validation of RP-HPLC Method for Simultenous Determination of cefixime and potassium Clavulanate in tablet dosage form. International Journal of Pharma Recent Research. 2010; 2(2): 57-60.
26. Ajit R.Wankhede, Prashant Y. Mali, Vikram Karne, Anubha R. Khale, C.S. Magdum. Development and validation of RP-HPLC method for simultenous estimation of cefixime and cloxacillin in tablet dosage form. International Journal of Pharmaceutical & Biological Archives. 2010; 1(2): 317-320.
27. Yost RL, Derendorf H., (1985), Rapid chromatographic determination of cefotaxime and its metabolite in biological fluids, J.Chromatogr., 341(1), 131-138.
28. Ling SS, Yuen KH., (2003), Simple liquid chromatographic method for the determination of cefotaxime in human and rat plasma. J. Chromatogr. B. Analyt. Technol. Biomed. Life Sci., 783(1), 297-301.
29. Martinez LG, Campins- Falco P, (1998), New spectrophotometric procedure for dtermining cefotaxime based on derivatization with 1,.2- naphtoquinone-4- sulphonate into solid phase extration cartridges. Application to pharmaceuticl and urine samples. J. Chromatogr. B. Biomed. Sci. Appi/. 718(1), 143-151.
30. Nuevas L, Gonzaiez R, (1998), Derivative spectrophotometric determination of the triethylammonium salt of cefotaxime in presence of related compounds from the synthesis. J. Pharm. and Biomed. Anal., 18(4-5), 579-583.
31. Scanes I, Hundt Af, Swart KJ, Jundt HK, (2001), simultaneous determination of cefotaxime and desacetylcefotaxime in human plasma and cerebrospinal fluid by HPLC. J.Chromatogr B Biomed Sci. Appl., 750(1), 171-176.
32. Dell D, Chamberlain J, Coppin F.J, (1981), Determination of cefotaxime and desacetylcefotaxime in plasma and urine by high-performance liquid chromatography, J Chromatogr., Dec 11;226(2):431-40.

33. Dokladalova J, Quercia GT, Stankewich JP, (1983), High-performance liquid chromatographic determination of cefoperazone in human serum and urine, J Chromatogr., Aug 12;276(1):129-37.
34. Li FS, Xu ZX, Xiao HB, Liang XM, (2000), Simultaneous determination of sulbactam sodium and cefoperazone sodium in sulperazon by high performance liquid chromatography, Se Pu., Nov;18(6):525-6.
35. Van Koutrik et al., (1999), Oral Dosage-Forms Containing a β- Lactam Antibiotics, United States Patent (Patent Number - 5,948,422), Sep. 7, 1999.
36. Flanner et al., (2012), Pharmaceutical Compositions and Methods for Improved Bacterial Eradication, United States Patent (Patent Number - US 8,299,052 B2), Oct. 30, 2012.
37. Yamaguchi et al., (2007), Beta-Lactam Antibiotic-Containing Tablet and Production Thereof and their Evolution Parameters. United States Patent (Patent Number - US 2007/0134.325 A1), Jun. 14, 2007.
38. Brans et al., (2014), Method for Measuring Beta-Lactam Antibiotics. United States Patent Application Publication (Publication Number - US 2014/0243241 A1), Aug. 28, 2014.
39. Terracciano et al., (2014), Oral Dosage-Forms Containing a β- Lactam Antibiotics their Method for Measuring Beta-Lactam Combinations. United States Patent Application Publication, United States Patent (Publication Number - US 2014/0274989 A1), Sep. 18, 2014.
40. N. H. Vadia, N. B. Dobaria and V. B. Patel, (2009), Conventional and Advanced Spectrophotometric Methods for estimation of Cefetamet Pivoxil Hydrochloride in Bulk and in Pharmaceutical Formulation, International Journal of ChemTech Research,Vol.1, No.2, pp 286-290, April-June 2009